

LATISSE® INFORMED CONSENT

1325 Warren Avenue Spring Lake NJ 08535 732 268-7663



Today's date ____/____/____ D O B ____/____/____

Patient name: _____

Address: _____
Street City State Zip

Home phone: _____ Cell: _____

E-mail _____

Referral _____

What is LATISSE®?

- LATISSE® is recommended for patients that have hypotrichosis (not enough eyelashes).
- LATISSE® solution is a prescription treatment for hypotrichosis used to grow eyelashes, making them longer, thicker and darker.
- LATISSE® solution is intended for use on the skin of the upper eyelid margins at the base of the eyelashes.

Contraindications:

- Pregnant women
- Patient allergic to any of LATISSE® ingredients and patients using LUMIGAN®.
- In patients using LUMIGAN® or other prostaglandin analogs for the treatment of elevated intraocular pressure, the concomitant use of LATISSE® may interfere with the desired reduction in IOP. Patients using prostaglandin analogs for IOP reduction should only use LATISSE® after consulting with their physician.
- Patients using contact lenses should be advised that LATISSE® solution contains benzalkonium chloride, which may be absorbed by soft contact lenses. Contact lenses should be removed prior to application of LATISSE® and may be reinserted 15 minutes following its administration.

Side effects:

- Potential for increase brown iris pigmentation which is likely to be permanent. Increased iris pigmentation has occurred when bimatoprost solution was administered.
- Possible hair growth occurring outside of the target treatment area if LATISSE® repeatedly touches the same area of skin outside the treatment area. There is also a possibility of disparity between eyes in length, thickness, pigmentation, number of eyelashes or vellus hairs, and/or direction of eyelash growth. Eyelash changes are likely reversible upon discontinuation of treatment.
- Possible eyelid skin darkening, which may be reversible after discontinuation of LATISSE®.
- LATISSE® may lower intraocular pressure although not to a level that will cause clinical harm.

How to apply LATISSE®

- LATISSE® (bimatoprost ophthalmic solution) should be applied every night using only the accompanying sterile applicators. They should start by ensuring their face is clean, all makeup is removed, and their contact lenses removed (if applicable). Then, carefully place one drop of LATISSE® solution on the disposable sterile applicator and brush cautiously along the skin of the upper eyelid margin at the base of the eyelashes. If any LATISSE® solution gets into the eye proper, it will not cause harm. The eye should not be rinsed.
- Additional applications of LATISSE® will not increase the growth of eyelashes.
- LATISSE® should not be applied to the lower eyelash line. Any excess solution outside the upper eyelid margin should be blotted with a tissue or other absorbent material.
- The onset of effect is gradual but is not significant in the majority of patients until 2 months. Patients should be counseled that the effect is not permanent and can be expected to gradually return to the original level upon discontinuation of treatment with LATISSE®.
- LATISSE® bottle must be maintained intact and to avoid allowing the tip of the bottle or applicator to contact surrounding structures, fingers, or any other unintended surface in order to avoid contamination of the bottle or applicator by common bacteria known to cause ocular infections.
- LATISSE® applicator supplied with the product should be used once and then discard since reuse could result in using a contaminated applicator. Serious infections may result from using contaminated solutions or applicators.

I _____ have read and understand all of the above and I aware of all the possible complications that may be caused by using LATISSE®, and I attest that I'm not allergic to any of the LATISSE® ingredients, I'm not pregnant and I'm not using LUMIGAN®.

Patient name _____

Patient signature _____